

ACIDE LACTIQUE OENO

CORRECTORS

90% lactic acid solution
Authorised dosage limit and acidification declaration.

↘ OENOLOGICAL APPLICATIONS

ACIDE LACTIQUE OENO is of natural origin, obtained by lactic sugar fermentation.

It is used to acidify musts derived from extremely ripe and insufficiently acidic grapes, or in wines.

ACIDE LACTIQUE OENO can be considered for usage before or after malolactic fermentation and in this case, it adds to the lactic acid content produced by the bacteria during malolactic fermentation.

↘ INSTRUCTIONS FOR USE

Dilute the **ACIDE LACTIQUE OENO** in 10 times its weight in must or wine, then incorporate it into the total volume to be treated.

↘ DOSE RATE

Dosage must be determined in the laboratory depending on total acidity, pH and tasting.

-1 g of TARTARIC ACID is equivalent to 1,1 mL of **ACIDE LACTIQUE OENO**

-1 g of H₂SO₄ is equivalent to 1,7 mL of **ACIDE LACTIQUE OENO**

- On white and rosé: 3 to 15 g/hL: for freshness and delicacy.
- On red: 10 to 20 g/hL: for freshness on the finish. Softens potential bitterness.

↘ PACKAGING AND STORAGE

- 1 L flask
- 10 L and 25 kg drum
- 1200 kg container

ACIDE LACTIQUE OENO must be stored in a dry, well-ventilated and odourless environment at a temperature between 5 and 25°C